

Growth index with the cosmological constant

Seokcheon Lee^{1,2} and Kin-Wang Ng^{1,2,3}

July 9, 2009

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, R.O.C.*

²*Leung Center for Cosmology and Particle Astrophysics, National Taiwan University, Taipei, Taiwan 10617, R.O.C.*

³*Institute of Astronomy and Astrophysics, Academia Sinica, Taipei, Taiwan 11529, R.O.C.*

Abstract

We obtain the exact analytic form of the growth index at present epoch ($a = 1$) in a flat universe with the cosmological constant (*i.e.* the dark energy with its equation of state $\omega_{de} = -1$). For the cosmological constant, we obtain the exact value of the current growth index parameter $\gamma = 0.5547$, which is very close to the well known value $6/11$. We also obtain the exact analytic solution of the growth factor for $\omega_{de} = -1/3$ or -1 . We investigate the growth index and its parameter at any epoch with this exact solution. In addition to this, we are able to find the exact Ω_m^0 dependence of those observable quantities. The growth index is quite sensitive to Ω_m^0 at $z = 0.15$, where we are able to use 2dF observation. If we adopt 2dF value of growth index, then we obtain the constrain $0.11 \leq \Omega_m^0 \leq 0.37$ for the cosmological constant model. Especially, the growth index is quite sensitive to Ω_m^0 around $z \leq 1$. We might be able to obtain interesting observations around this epoch. Thus, the analytic solution for this growth factor provides the very useful tools for future observations to constrain the exact values of observational quantities at any epoch related to growth factor for $\omega_{de} = -1$ or $-1/3$.

The background evolution equations in a flat Friedmann-Robertson-Walker universe ($\rho_m + \rho_{de} = \rho_{cr}$) are

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}(\rho_m + \rho_{de}) = \frac{8\pi G}{3}\rho_{cr}, \quad (1)$$

$$2\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2 = -8\pi G\omega_{de}\rho_{de}, \quad (2)$$

where ω_{de} is the equation of state (eos) of dark energy, ρ_m and ρ_{de} are the energy densities of the matter and the dark energy, respectively. We consider the constant ω_{de} only. The dark energy does not participate directly in cluster formation, but it alters the cosmic evolution of the background. The linear density perturbation of the matter ($\delta = \delta\rho_m/\rho_m$) for sub-horizon scales is governed by [1]

$$\ddot{\delta} + 2\frac{\dot{a}}{a}\dot{\delta} = 4\pi G\rho_m\delta. \quad (3)$$

The textbook solution for the growing mode solution of Eq. (3) for $\omega_{de} = -1$ or $-1/3$ is [2, 3, 4]

$$\delta_g(a) = \frac{5\Omega_m^0}{2} \frac{H(a)}{H_0} \int_0^a X^{-3/2}(a') da', \quad (4)$$

where $X(a) = (aH/H_0)^2 = \Omega_m^0 a^{-1} + \Omega_{de}^0 a^{-1-3\omega_{de}}$, $\Omega_m^0 = \rho_m^0/\rho_{cr}^0$ and $\Omega_{de}^0 = 1 - \Omega_m^0$ are the present energy density contrast of the matter and the dark energy, respectively. The growth index f is defined as

$$f = \frac{d \ln \delta_g}{d \ln a} \equiv \Omega_m(a)^\gamma. \quad (5)$$

Thus, f is expressed from Eq. (4) as

$$f(\Omega_m^0, \omega_{de}, a) = -\frac{3}{2} - \frac{3\omega_{de}(1 - \Omega_m^0)a^{-1-3\omega_{de}}}{2X} + \frac{aX^{-3/2}(a)}{\int_0^a X^{-3/2}(a') da'}. \quad (6)$$

We obtain the exact analytic form of the growth index for $\omega_{de} = -1$ or $-1/3$ at the present epoch ($a = 1$, $X(a = 1) = 1$) as

$$f(\Omega_m^0, \omega_{de}, 1) = -\frac{3}{2} - \frac{3}{2}\omega_{de}(1 - \Omega_m^0) - 3\omega_{de}(\Omega_m^0)^{\frac{3}{2}} \frac{\Gamma[1 - \frac{5}{6\omega_{de}}]/(\Gamma[\frac{-5}{6\omega_{de}}]\Gamma[1])}{F[\frac{3}{2}, \frac{-5}{6\omega_{de}}, 1 - \frac{5}{6\omega_{de}}, -\frac{\Omega_{de}^0}{\Omega_m^0}]}, \quad (7)$$

where Γ is the gamma function and F is the hypergeometric function. We emphasize that this formula is correct only for $\omega_{de} = -1/3$ or -1 . For $\Omega_m^0 = 1$ one get $f = 1$ for all a , which is consistent with $\delta \propto a$. For $\omega_{de} = -\frac{1}{3}$, f is matched with a non-flat without the cosmological constant $f(a = 1) = -1 - \Omega_m^0/2 + 5/2(\Omega_m^0)^{3/2}/F[3/2, 5/2, 7/2, 1 - (\Omega_m^0)^{-1}]$ as shown in Ref. [5]. For the cosmological constant, the above formula becomes

$$f(\Omega_m^0, -1, 1) \equiv f_L^0 = -\frac{3}{2}\Omega_m^0 + 3(\Omega_m^0)^{\frac{3}{2}} \frac{\Gamma[\frac{11}{6}]/\Gamma[\frac{5}{6}]}{F[\frac{3}{2}, \frac{5}{6}, \frac{11}{6}, -\frac{\Omega_{de}^0}{\Omega_m^0}]}, \quad (8)$$




Figure 1: Ω_m^0 dependence of f_L^0 and γ_L^0 . a) Current value of the growth index f_L^0 for $0.2 \leq \Omega_m^0 \leq 0.4$. b) γ_L^0 for the same range of Ω_m^0 .

where we use $\Gamma[1] = 1$. From the exact form of the growth index f_L^0 , we are able to obtain the exact present value of the growth index parameter γ_L^0 ,

$$\gamma_L^0 \equiv \frac{\ln f_L^0}{\ln \Omega_m^0} = \frac{\ln \left[-\frac{3}{2} \Omega_m^0 + (\Omega_m^0)^{\frac{3}{2}} \frac{\Gamma[\frac{11}{6}]/\Gamma[\frac{5}{6}]}{F[\frac{3}{2}, \frac{5}{6}, \frac{11}{6}, -\Omega_{de}^0/\Omega_m^0]} \right]}{\ln \Omega_m^0}. \quad (9)$$

In Fig. 1, we show the present value of the growth index f_L^0 and its parameter γ_L^0 as a function of Ω_m^0 . In Fig. 1a, we show the variation of the present growth index for different values of Ω_m^0 . It is changed from 0.4073 to 0.6028 for $0.2 \leq \Omega_m^0 \leq 0.4$. Thus, there is up to 32% differences in f_L^0 for the different values of Ω_m^0 . However, this Ω_m^0 dependence is decreased by using the parameter γ_L^0 . We show this variation of γ_L^0 in Fig. 1b. γ_L^0 is changed from 0.5580 to 0.5524 for $\Omega_m^0 = 0.2$ and 0.4, respectively. We obtain $\gamma_L^0 = 0.5547$ for $\Omega_m^0 = 0.3$.

We also obtain the exact analytic solution of the growth factor given in Eq. (4) for $\omega_{de} = -1/3, -1$ (see Appendix for details). The solution of Eq. (16) for $\omega_{de} = -1$ is

$$\delta_g^L(a) = c_1^L \sqrt{1 + Qa^{-3}} + c_2^L Q^{\frac{2}{3}} a^{-2} F\left[1, \frac{1}{6}, \frac{5}{3}, -Qa^{-3}\right], \quad (10)$$

where $c_{1,2}^L$ are integral constants and $Q = \Omega_m^0/\Omega_{de}^0$. We need to determine the c_1^L and c_2^L based on the proper initial conditions $\delta_g^L(a_i) = a_i$ and




Figure 2: Cosmological evolution of δ_g^L for $\Omega_m^0 = 0.4, 0.3, 0.2$ (from top to bottom).

$d\delta_g^L/da|_{a_i} = 1$ in the matter dominated epoch in order to make this solution become the growing mode solution. We obtain from the initial conditions that $c_1^L = 1.085464$ and $c_2^L = -0.943314$ when $\omega_{de} = -1.0$ and $\Omega_m^0 = 0.3$.

We show the cosmological evolution of the growth factor δ_g^L for different values of Ω_m^0 in Fig. 2. The dotted, solid, and dashed lines correspond to $\Omega_m^0 = 0.4, 0.3$ and 0.2 respectively. As Ω_m^0 increases, δ_g^L maintains the linear growth with a for a longer time, as expected. From the initial conditions, we obtain $(c_1^L, c_2^L) = (1.25633, -1.09263)$ and $(0.906875, -0.788711)$ for $\Omega_m^0 = 0.4$ and 0.2 , respectively. We use these values of the coefficients in Fig. 2.

In addition to the present value of the growth index $f(a = 1)$, we are able to obtain the growth index in any epoch from the exact analytic solution of the growth factor (10). For $\omega_{de} = -1$, we find that

$$f_L(a) = \frac{d \ln \delta_g^L(a)}{d \ln a} = \frac{\left(-\frac{3}{2} \frac{A[a]^2 - (c_1^L)^2}{A[a]} - 2B[a] + \frac{3}{10} Q a^{-3} B[a] \frac{F2}{F1} \right)}{(A[a] + B[a])}$$

where,

$$A[a] = c_1^L \sqrt{1 + Q a^{-3}}, \quad B[a] = c_2^L Q^{\frac{3}{2}} a^{-2} F1 \quad (11)$$

$$F1 = F\left[1, \frac{1}{6}, \frac{5}{3}, -Q a^{-3}\right], \quad F2 = F\left[2, \frac{7}{6}, \frac{8}{3}, -Q a^{-3}\right]$$




Figure 3: a) Evolution of $f_L(a)$ for different values of Ω_m^0 . b) The same for $\gamma_L(a)$. Dotted, solid, and dashed lines correspond to $\Omega_m^0 = 0.4, 0.3$, and 0.2 , respectively.

Even though it looks complicated, one is able to obtain $f_L(a)$ by using the solution δ_g^L without any difficulty. We are able to find the growth index parameter $\gamma_L(a) = \frac{\ln f_L(a)}{\ln \Omega_m(a)}$ in any epoch from this exact analytic form of $f_L(a)$. We also investigate the Ω_m^0 dependence of them without any ambiguity. Thus, this analytic solution is very useful for the investigation of observational quantities. We show these properties in Fig. 3.

In Fig. 3a, we show the growth index $f_L(a)$ for different values of Ω_m^0 . The dotted, solid and dashed lines correspond to the evolution of $f_L(a)$ for $\Omega_m^0 = 0.4, 0.3$ and 0.2 , respectively. As we have more matter at present, we have faster growing. Thus, we have bigger values of f_L when we have bigger values of Ω_m^0 as shown in Fig. 3a. The Ω_m^0 dependence is quite sensitive around $a \simeq 0.87$ (*i.e.* $z \simeq 0.15$). Thus, the observed value of $f(a = 0.87) = 0.51 \pm 0.15$ from the 2dF galaxy redshift survey will be a good guideline to measure Ω_m^0 if the dark energy is the cosmological constant [6]. However, the present value of the growth index parameter is insensitive to Ω_m^0 as we also obtain in the previous formula. The growth index parameter is sensitive to Ω_m^0 around $a \geq 0.8$ (*i.e.* $z \leq 0.25$) as shown in Fig. 3b. Again, the dashed, solid and dotted lines correspond to the cosmological evolution of $\gamma_L(a)$ for $\Omega_m^0 = 0.2, 0.3$ and 0.4 , respectively. Especially, the growth index




Figure 4: Cosmological evolution of $\gamma_L(a)$ for $\Omega_m^0 = 0.2, 0.3, 0.4$ (from top to bottom).

parameter changes dramatically for $a \leq 0.3$ (*i.e.* $z \geq 2.3$). We show this in Fig. 4. The dashed, solid and dotted lines (from top to bottom) correspond to the evolution of $\gamma_L(a)$ for $\Omega_m^0 = 0.2, 0.3$ and 0.4 , respectively.

We summarize the results in Table 1. As it is well known that the growth index parameter γ_L is insensitive to Ω_m^0 and a up to $a > 0.3$ [7]. However, $\gamma_L(a)$ shows the strong model dependence around $a \leq 0.3$ (*i.e.* $z \geq 2.3$). If we naively adopt the 2dF value of f without considering the data error from selection effects, then we obtain the constrain $0.11 \leq \Omega_m^0 \leq 0.37$ for the cosmological constant model.

In linear theory, the peculiar velocity $\vec{v}_{pec}$ is related to the peculiar acceleration $\vec{g}$ and/or interior average over-density $\langle \delta \rangle_R$ in a spherical perturbation of radius R , [8]

$$|\vec{v}_{pec}| = \frac{2}{3} \left| \frac{f\vec{g}}{H\Omega_m^0} \right| = \frac{1}{3} H R f \langle \delta \rangle_R. \quad (12)$$

The difference of this peculiar velocity between two different values of Ω_m^0 is as large as 29% when we use the different values of $f_L^{z=0.15}$ for $\Omega_m^0 = 0.2$ and 0.4 . Thus, the exact analytic form of f provides the good analysis tool for galaxy redshift survey.

Ω_m^0	$f_L^{z=0}$	$\gamma_L^{z=0}$	$f_L^{z=0.15}$	$\gamma_L^{z=0.15}$
0.2	0.407	0.558	0.488	0.555
0.3	0.513	0.555	0.598	0.553
0.4	0.603	0.552	0.685	0.551

Table 1: Ω_m^0 is the present value of the matter density contrast, $f_L^{z=0}$ and $f_L^{z=0.15}$ correspond to the growth index at the present and $z = 0.15$, respectively. $\gamma_L^{z=0}$ and $\gamma_L^{z=0.15}$ are the growth index parameter at the present and $z = 0.15$, respectively.

A Appendix

From Eq. (6), we need to solve the integration to find the analytic form of f at the present epoch [10],

$$\begin{aligned}
\int_0^1 \frac{da}{X^{3/2}(a)} &= \int_0^1 \frac{da}{(\Omega_m^0 a^{-1} + \Omega_{de}^0 a^{-1-3\omega_{de}})^{3/2}} = \int_0^1 \frac{da}{(1 + \Omega_{de}^0/\Omega_m^0 a^{-3\omega_{de}})^{3/2} (\Omega_m^0/a)^{3/2}} \\
&= \frac{-1}{3\omega_{de}(\Omega_m^0)^{3/2}} \int_0^1 \eta^{-1-5/(6\omega_{de})} (1 - r\eta)^{-3/2} d\eta \\
&= \frac{-1}{3\omega_{de}(\Omega_m^0)^{3/2}} \frac{\Gamma[\frac{-5}{6\omega_{de}}]\Gamma[1]}{\Gamma[1 - \frac{5}{6\omega_{de}}]} F\left[\frac{3}{2}, \frac{-5}{6\omega_{de}}, 1 - \frac{5}{6\omega_{de}}, -\frac{\Omega_{de}^0}{\Omega_m^0}\right], \tag{13}
\end{aligned}$$

where we use $a^{-3\omega_{de}} = \eta$ and $r = -\Omega_{de}^0/\Omega_m^0 = 1 - (\Omega_m^0)^{-1}$. Note that Eq. (13) is valid for any value of ω_{de} . We can also check the result of a non-flat universe without the cosmological constant in Ref. [5]. In this case we have $k/H_0^2 \equiv \Omega_K = 1 - \Omega_m^0$. Then we obtain the expression for $X = \Omega_m^0 a^{-1} + \Omega_K$. Mathematically, this is the same as the dark energy case with the equation of state of the dark energy $\omega_{de} = -1/3$. Then the above integral (13) becomes

$$\int_0^1 \frac{da}{X^{3/2}(a)} = \frac{1}{(\Omega_m^0)^{3/2}} \frac{\Gamma[\frac{5}{2}]\Gamma[1]}{\Gamma[\frac{7}{2}]} F\left[\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, -\frac{\Omega_K}{\Omega_m^0}\right] = \frac{2}{5} (\Omega_m^0)^{-3/2} F\left[\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, 1 - (\Omega_m^0)^{-1}\right]. \tag{14}$$

Thus, we can reproduce $f = -1 - \Omega_m^0/2 + 5/2(\Omega_m^0)^{3/2}/F[\frac{3}{2}, \frac{5}{2}, \frac{7}{2}, 1 - (\Omega_m^0)^{-1}]$ in a non-flat without the cosmological constant case.

It is better for us to rewrite the linear density perturbation equation (3) with changing the variable from t to a to get [4]

$$\frac{d^2\delta}{da^2} + \left(\frac{d\ln H}{da} + \frac{3}{a}\right) \frac{d\delta}{da} - \frac{4\pi G\rho_m}{(aH)^2} \delta = 0. \tag{15}$$

When we replace the growing mode solution (4) into Eq. (3), we get

$$\ddot{\delta}_g^{ex} + 2H\dot{\delta}_g^{ex} - 4\pi G\rho_m\delta_g^{ex} - \left[4\pi G(1 + \omega_{de})(1 + 3\omega_{de})\rho_{de}\right]\delta_g^{ex} = 0. \quad (16)$$

Note that we have changed the notation from δ_g to δ_g^{ex} to reflect the fact that δ_g^{ex} is the solution of Eq. (16) for any value of ω_{de} . In particular, when $\omega_{de} = -1$ or $-1/3$, the extended solution δ_g^{ex} is reduced to δ_g . After replacing new parameters $Y = (\Omega_m)/(\Omega_{de}) = (\Omega_m^0)/(\Omega_{de}^0)a^{3\omega_{de}}$ in Eq. (16), we have

$$Y\frac{d^2\delta_g^{ex}}{dY^2} + \left[1 + \frac{1}{6\omega_{de}} - \frac{1}{2(Y+1)}\right]\frac{d\delta_g^{ex}}{dY} - \left[\frac{1}{6\omega_{de}^2 Y} + \frac{3\omega_{de} + 4}{6\omega_{de}Y(Y+1)}\right]\delta_g^{ex} = 0. \quad (17)$$

Now we try $\delta_g^{ex}(Y) = c_1\delta_1(Y) + c_2\delta_2(Y) = c_1Y^i(1+Y)^j + c_2Y^k B(Y)$, where $c_{1,2}$ are integral constants, because it is the most general combination of the solution for the above equation (17). We obtain the constraints for i and j from δ_1 ,

$$i = -\frac{1 + \omega_{de}}{2\omega_{de}}, \quad j = \frac{1}{2}. \quad (18)$$

We also replace δ_2 into the above equation (17) to get

$$Y(1+Y)\frac{d^2B}{dY^2} + \left[\frac{5}{6\omega_{de}} + \frac{5}{2} + \left(3 + \frac{5}{6\omega_{de}}\right)Y\right]\frac{dB}{dY} + \left(1 + \frac{5}{6\omega_{de}}\right)B = 0, \\ \text{when } k = 1 + \frac{1}{3\omega_{de}}. \quad (19)$$

There are two alternative ways to make the above equation as the hypergeometric differential equation, $Y = -X$ or $1 + Y = X$ [9]. We choose the first case which shows the proper behavior. Now the above equation (19) becomes the so-called hypergeometric differential equation,

$$(X)(1-X)\frac{d^2B}{dX^2} + \left[\frac{1}{2} - \frac{18\omega_{de} + 5}{6\omega_{de}}X\right]\frac{dB}{dX} - \left(1 + \frac{5}{6\omega_{de}}\right)B = 0, \quad (20)$$

with the solution $B(X) = F\left[1, 1 + \frac{5}{6\omega_{de}}, \frac{5}{2} + \frac{5}{6\omega_{de}}, X\right]$ being the hypergeometric function [9]. Thus, the full extended solution becomes

$$\delta_g^{ex}(Y) = c_1Y^{-\frac{1+\omega_{de}}{2\omega_{de}}}\sqrt{1+Y} + c_2Y^{\frac{1+3\omega_{de}}{3\omega_{de}}}F\left[1, 1 + \frac{5}{6\omega_{de}}, \frac{5}{2} + \frac{5}{6\omega_{de}}, -Y\right]. \quad (21)$$

Writing $Y = a^{3\omega_{de}}Q$, $Q = \Omega_m^0/\Omega_{de}^0$, for $\omega_{de} = -1$ and $-1/3$, we have

$$\begin{aligned}\delta_g^{ex}(a)_{\omega_{de}=-1} &= \delta_g^L(a) = c_1^L \sqrt{1 + Qa^{-3}} + c_2^L Q^{\frac{2}{3}} a^{-2} F\left[1, \frac{1}{6}, \frac{5}{3}, -Qa^{-3}\right] \\ \delta_g^{ex}(a)_{\omega_{de}=-\frac{1}{3}} &= c_1 Qa^{-1} \sqrt{1 + Qa^{-1}} + c_2 F\left[1, -\frac{2}{3}, 0, -Qa^{-1}\right].\end{aligned}\quad (23)$$

The solution (21) is a mathematical one. We need to further specify this solution in order to make it as a physical solution. If we use the fact that growth factor $\delta \propto a$ in the matter dominated epoch, then we can determine c_1 and c_2 to make the solution as the growing mode solution. We are also able to find the decaying mode solution if we use the decaying mode initial conditions into this solution (21).

References

- [1] W. B. Bonnor, Mon. Not. R. Astron. Soc. **117**, 104 (1957).
- [2] D. J. Heath, Mon. Not. R. Astron. Soc. **179**, 351 (1977).
- [3] V. Sahni and P. Coles, Phys. Rep. **262**, 1 (1995) [arXiv:astro-ph/9505005]; U. Alam, V. Sahni, and A. A. Starobinsky, [arXiv:0812.2846].
- [4] S. Dodelson, *Modern Cosmology* (Academic Press, San Diego, 2002).
- [5] O. Lahav, P. B. Lilje, J. R. Primack, and M. J. Rees, Mon. Not. R. Astron. Soc. **251**, 128 (1991).
- [6] E. Hawkins *et al.*, Mon. Not. R. Astron. Soc. **346**, 78 (2003) [arXiv:astro-ph/0212375], L. Verde *et al.*, Mon. Not. R. Astron. Soc. **335**, 432 (2002) [arXiv:astro-ph/0112161].
- [7] L. Wang and P. J. Steinhardt, Astrophys. J. **508**, 483 (1998) [astro-ph/9804015].
- [8] P. J. E. Peebles, *Large Scale Structure of the Universe* (Princeton University Press, Princeton, 1980).
- [9] P. M. Morse and H. Feshbach, *Methods of Theoretical Physics, Part I* (McGraw-Hill Science, New York, 1953).

- [10] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables* (Dover Publications, New York, 1970).